



Extension

UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

Late Blight

Amanda Gevens, Anna Seidl Johnson and Brian Hudelson, UW-Madison Plant Pathology

What is late blight? Late blight is a destructive disease of tomatoes and potatoes that can kill plants, and make tomato fruits and potato tubers inedible. This disease also affects, although typically to a lesser extent, weeds such as nightshade. Historically, late blight was responsible, in part, for the Irish potato famine, which led to the death of at least one million Irish, and the emigration of another one million Irish to Great Britain, the United States, Australia, Canada and other countries. The disease routinely causes severe losses for fresh market tomato producers and home gardeners in Wisconsin.



Late blight can decimate tomato and potato plants in seven to 10 days if weather conditions are cool and wet.

What does late blight look like?

On leaves of tomato or potato, late blight begins as pale-green or olive-green areas that quickly enlarge to become brown-black, water-soaked, and oily-looking. Stems can also exhibit dark-brown to black areas. If weather conditions are cool and wet, entire plants can collapse and die from late blight in seven to 10 days. Tomato fruits with late blight develop large, often sunken, golden- to chocolate-brown, firm spots with distinct rings. Potato tubers with the disease develop a reddish-brown discoloration under the skin and these areas may become sunken. Affected

leaf, stem, fruit or tuber tissue often eventually develops a white-gray, fuzzy look as the late blight organism begins to reproduce. Other tomato and potato diseases such as Septoria leaf spot (see UW Plant Disease Facts D0100, *Septoria Leaf Spot*), early blight (see UW Plant Disease Facts D0046, *Early Blight*) and Verticillium wilt (see UW Plant Disease Facts D0122, *Verticillium Wilt of Vegetables*) may be misidentified as late blight.

Where does late blight come from? Late blight is caused by the fungus-like water mold *Phytophthora infestans*. There are several variants or strains of this organism. Some variants/strains cause more severe problems on tomatoes; others are more of a problem on potatoes. *P. infestans* can be introduced when sporangia (i.e., spore-like structures) of the organism are blown into an area on prevailing winds. *P. infestans* can also be introduced on infected plants (e.g., tomato seedlings) or potato tubers that are brought into an area for sale. *P. infestans* is unlikely to survive over the winter in Wisconsin soils. However, the organism can survive on infected plant material that remains alive through the winter. Such plant materials can include *P. infestans*-infected tomato plants or fruits kept warm in compost piles, and *P. infestans* infected potato tubers that have remained in the soil after harvest or have been stored in a warm place.

How do I save a plant with late blight? Plants showing symptoms of late blight cannot be saved and should be disposed of immediately to limit spread of *P. infestans* to other plants. For home gardeners, the preferred method of disposal is to pull affected plants (roots and all), and place them in plastic bags. The bags should be left in the sun for a few days to make sure that plants, as well as any *P. infestans*, are totally killed. Bagged plants should then be put out for trash pickup. Diseased plants or plant parts (e.g., tomato fruits or potato tubers) should NOT be composted. Healthy-looking fruits from late-blight-affected tomato plants are safe to eat or preserve. However, before attempting to preserve any fruit or vegetable, contact your county UW-Extension office for the most current food preservation instructions. Once tomato fruits or potato tubers begin to show symptoms of late blight, they should NOT be eaten, nor should they be canned or otherwise preserved.



Extension
UNIVERSITY OF WISCONSIN-MADISON



On tomato leaves (left), late blight leads to brown-black, water-soaked, oily areas that may have a white-gray fuzzy look. On tomato fruits (right), late blight leads to large, often sunken, golden- to chocolate-brown, firm spots with distinct rings.

How do I avoid problems with late blight in the future? In the spring, dispose of any volunteer tomato and potato plants (as well as weeds such as nightshade) as described above. All of these plants are potential sources of *P. infestans*. For the same reason, DO NOT use tubers from a previous year's potato crop as seed potatoes. Instead purchase certified seed potatoes from a reputable supplier each year. Also, consider planting tomato varieties with late blight tolerance or resistance. Such varieties include 'Defiant PHR', 'Iron Lady', 'Jasper', 'JTO-545', 'Lemon Drop', 'Matt's Wild Cherry', 'Mountain Magic', 'Mountain Merit', 'Mr. Stripey', 'Plum Regal', 'Pruden's Purple', 'Stellar' and 'Wapsipinicon Peach.' The performance of these varieties may vary depending on the variant or strain of *P. infestans* that is present in a particular growing season, and depending on the weather conditions. Consider planting several resistant tomato varieties to help ensure that no matter what *P. infestans* variant is prevalent in a given year, you will have some healthy tomato plants and fruits.

Fungicides also can be used to reduce the impact of late blight. However, fungicide applications must be made prior to the onset of disease or they will be ineffective. Fungicide applications are not needed during periods of hot, dry weather as *P. infestans* is not likely to be active under these conditions. Fungicides are most likely to be useful during periods of cool, wet weather. However, if weather conditions are excessively cool and wet, even properly-timed fungicide applications may not provide adequate late blight control. If you decide to use fungicides, select a product that is labeled for use on tomatoes (or potatoes) and that contains chlorothalonil or copper as the active ingredient. Certain, but not all, copper-containing products can be used for organic vegetable production. Be sure to follow all label instructions to ensure that you use the product that you select in the safest, most effective means possible. For more information on home garden fungicides for vegetable disease control, see UW Plant Disease Facts D0062, *Home Vegetable Garden Fungicides*.

For more information on late blight: Contact the University of Wisconsin Plant Disease Diagnostics Clinic (PDDC) at (608) 262-2863 or pddc@wisc.edu.

© 2010-2024 by the Board of Regents of the University of Wisconsin System doing business as the Division of Extension of the University of Wisconsin-Madison.

An EEO/Affirmative Action employer, University of Wisconsin-Madison provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Kristin Krokowski, Barb Larson, Patti Nagai and Ann Wied for reviewing this document.

A complete inventory of UW Plant Disease Facts is available at the University of Wisconsin-Madison Plant Disease Diagnostics Clinic website: <https://pddc.wisc.edu>.

Submit additional lawn, landscape, and gardening questions at <https://hort.extension.wisc.edu/ask-a-gardening-question/>.